

## Journal of Pharmaceutical Advanced Research

(An International Multidisciplinary Peer Review Open Access monthly Journal)

Available online at: [www.jparonline.com](http://www.jparonline.com)**A review on Taurine - A magic molecule**Simanchal Panda<sup>1\*</sup>, Sruti Ranjan Mishra<sup>2</sup>, V.V.V.K. Mishra<sup>2</sup><sup>1</sup>M.D.A.M Doctorate of Alternative Medicine, P.I., IFS.<sup>2</sup>Jeypore College of Pharmacy, Jeypore – 764002, Koraput, Odisha.

Received: 26.02.2018

Revised: 18.03.2018

Accepted: 22.03.2018

Published: 31.03.2018

**ABSTRACT:** Taurine is an amino sulfonic acid (Amino acid), a chemical that is a required building block of protein. Taurine is found in large amounts in the brain, retina, heart, and blood platelets. The best food sources are meat and fish. Early research suggests that taking a nutritional supplement containing taurine improves vision in eye of people with age-related macular degeneration (AMD). The research suggests that taking taurine improves the symptoms of nausea and vomiting in patients receiving chemotherapy. Taurine supplementation might be useful along with usual treatment to reduce fatty stools (steatorrhea) in children with cystic fibrosis. The study suggests that taking 1.5 g of taurine twice daily for 4 months does not affect blood sugar, blood fats, or insulin levels in people with diabetes. Daily energy drink containing taurine prior to driving may reduce driver fatigue. About 1 to 1.66 g of taurine before exercise does not improve overall exercise performance. Using products containing taurine combined with other ingredients might improve cycling performance but not strength training or sprint performance. About 500 mg of taurine twice daily together with conventional treatments for 6 weeks reduces stomach ulcers caused by *Helicobacter pylori* infection. Taurine give therapeutic benefits for high blood pressure, anaemia, mental performance and muscle soreness. The research suggests that taking 100-150 mg/kg of taurine for 6 months improves the ability to relax muscles after use in people with myotonic dystrophy and lack of sleep in combination with caffeine for sleep deprived patient.

**Corresponding author\***

Mr. Simanchal Panda  
M.D.A.M Doctorate of Alternative Medicine,  
P.I., IFS, India.  
E. Mail ID. [phrpvtltd@gmail.com](mailto:phrpvtltd@gmail.com)  
Tel. No. +919090870494.

**INTRODUCTIONS:**

Taurine, 2-aminoethanesulfonic acid is an organic compound that is widely distributed in animal tissues [1]. It is a major constituent of bile and can be found in the large intestine, and accounts for up to 0.1 % of total human body weight. Taurine has many fundamental biological roles, such as conjugation of bile acids, anti-oxidation, osmoregulation, membrane stabilization and modulation of calcium signalling. It is essential for cardiovascular function, and development and function of skeletal muscle, the retina, and the central nervous system. Taurine is unusual among biological molecules

**Key words:** Amino sulfonic acid, Cystic fibrosis, Myotonic dystrophy, Neuroprotection, Thermoregulation, Retinal protection, Seizures.

in being a sulfonic acid, while the vast majority of biologically occurring acids contain the more weakly acidic carboxyl group. While taurine is sometimes called an amino acid, and indeed is an acid containing an amino group, it is not an amino acid in the usual biochemical meaning of the term, which refers to compounds containing both an amino and a carboxyl group<sup>[2]</sup>.

#### CHEMISTRY OF TAURINE:

Taurine is 2-aminoethanesulfonic acid. Its molecular formula is Chemical formula  $C_2H_7NO_3S$ . The Molecular Mass is 125.14 g/mol. Its density is 1.734 g/cm<sup>3</sup>. Taurine possess melting point of 305.11 °C (581.20 °F; 578.26 K). Taurine is named after the Latin taurus (a cognate of the Greek τaurus) which means bull or ox, as it was first isolated from ox bile in 1827 by German scientists Friedrich Tiedemann and Leopold Gmelin<sup>[3]</sup>.

#### BIOSYNTHESIS OF TAURINE:

Taurine is derived from cysteine. Mammalian taurine synthesis occurs in the pancreas via the cysteine sulfinic acid pathway. In this pathway, cysteine is first oxidized to its sulfinic acid, catalyzed by the enzyme cysteine dioxygenase. Cysteine sulfinic acid, in turn, is decarboxylated by sulfinoalanine decarboxylase to form hypotaurine. Hypotaurine is enzymatically oxidized to yield taurine by hypotaurine dehydrogenase<sup>[4]</sup>. Taurine is also produced by the transsulfuration pathway, which converts homocysteine into cystathionine. The cystathionine is then converted to hypotaurine by the sequential action of three enzymes: cystathionine gamma-lyase, cysteine dioxygenase, and cysteine sulfinic acid decarboxylase. Hypotaurine is then oxidized to taurine.

#### Oxidative degradation of cysteine to taurine:

An important function of the  $\alpha$ -amino acid, taurine is the regulation of oxidative stress. However, taurine is neither a classical scavenger nor a regulator of the antioxidative defenses, leaving uncertain the mechanism underlying the antioxidant activity of taurine. In the present study, the taurine antagonist and taurine transport inhibitor,  $\beta$ -alanine, was used to examine the mechanism underlying the antioxidant activity of taurine. Exposure of isolated cardiomyocytes to medium containing  $\beta$ -alanine for a period of 48 h led to a 45 % decrease in taurine content and an increase in mitochondrial oxidative stress, as evidenced by enhanced superoxide generation, the inactivation of the

oxidant sensitive enzyme, aconitase, and the oxidation of glutathione. Associated with the increase in oxidative stress was a decline in electron transport activity, with the activities of respiratory chain complexes I and III declining 50-65 % and oxygen consumption falling 30%. A reduction in respiratory chain activity coupled with an increase in oxidative stress is commonly caused by the development of a bottleneck in electron transport that leads to the diversion of electrons from the respiratory chain to the acceptor oxygen forming in the process superoxide. Because  $\beta$ -alanine exposure significantly reduces the levels of respiratory chain complex subunits, ND5 and ND6, the bottleneck in electron transport appears to be caused by impaired synthesis of key subunits of the electron transport chain complexes. Co-administration of taurine with  $\beta$ -alanine largely prevents the mitochondrial effects of  $\beta$ -alanine, but treatment of the cells with 5 mM taurine in the absence of  $\beta$ -alanine has no effect on the mitochondria, likely because taurine treatment has little effect on cellular taurine levels. Taurine act as mitochondrial protein synthesis regulator, thereby enhancing electron transport chain activity and protecting the mitochondria against excessive superoxide generation<sup>[5-7]</sup>.

#### FUNCTIONS OF TAURINE:

Taurine was first isolated from ox bile, and thus its name is derived from the Latin 'Taurus', meaning 'ox' or 'bull'. Taurine is an amino acid and is present extensively in animal tissue. It is found in meat and fish, human tissue, large intestine and human breast milk. Taurine helps to improve blood pressure, cardiovascular function, retina, skeletal muscles. Taurine is usually added in energy drinks, infant formula, contact lens solutions, and cosmetics. Yes, taurine is present in bull sperm but that is not its source for energy drinks. Like many other additives, taurine too is synthesized in lab. It is manufactured by chemical synthesis in food processing laboratory<sup>[8,9]</sup>.

#### SIDE EFFECTS OF TAURINE:

The main negative side effect associated with taurine supplementation is negative nitrogen balance. The University of Utah Health Care Center reports that consuming a single amino acid supplement like taurine can offset your natural nitrogen balance. This may lead to a decrease in your metabolic efficiency and a strain on your kidneys. Children taking taurine supplements may also experience stunted growth<sup>[10]</sup>.

**DRUG INTERACTIONS:**

Research into drug interactions with taurine is sparse and inconclusive, according to the University of California San Diego. Some research points to chemotherapy drugs like cisplatin and fluorouracil affecting how well your body absorbs taurine. Though the long term and drug contraindications for taurine are not well known, you should still speak to your physician about possible complications from taurine supplementation before taking products that contain the amino acid [11].

**WHY TAURINE IS WONDER MOLECULE?**

Early research suggests that taking a nutritional supplement containing taurine by mouth, in addition to standard care for 6 months, improves the vision in people with Age mediated Diseases. Early research suggests that taking taurine by mouth improves the symptoms of nausea in vomiting in patients receiving chemotherapy. Taurine supplementation might be useful along with usual treatment to reduce fatty stools (steatorrhea) in children with cystic fibrosis. However, it does not seem to improve growth, lung function, or other symptoms of cystic fibrosis. Early research suggests that taking 1.5 g of taurine twice daily for 4 months does not affect blood sugar, blood fats, or insulin levels in people with diabetes. The scientific study suggests that taking an energy drink containing taurine prior to driving may reduce driver fatigue. Research suggests that taking 1-1.66 g of taurine before exercise does not improve overall exercise performance. Using products containing taurine combined with other ingredients might improve cycling performance but not strength training or sprint performance. Stomach ulcers caused by *H. pylori* infection being cured by taking 500 mg of taurine twice daily together with conventional treatments for 6 weeks. The extensive research suggests that taking 6 g of taurine daily for 7 days reduces blood pressure in people with borderline high blood pressure. The research suggests that taking iron with 1000 mg of taurine improves red blood cell counts and iron levels in women with anaemia due to iron deficiency. Early clinical research suggests that taurine, in combination with caffeine and Vitamin B complex, can improve attention and reasoning in adolescents, but does not improve memory. Research suggests that taking 2 g of taurine together with branched chain amino acids three times daily for 2 weeks reduces muscle soreness in healthy people who do not exercise regularly. Early research

suggests that taking 100-150 mg/kg of taurine for 6 months improves the ability to relax muscles after use in people with inherited muscle wasting disease that is myotonic dystrophy. The study suggests that taking taurine plus caffeine or a combination product containing taurine, caffeine, and vitamins B complex, reduces sleepiness and improves reaction time in people who are sleep deprived [10-15].

**CONCLUSION:**

In the present review, attempts have been made to provide synopsis, synthesis and symbiosis of chemical and biological actions of Taurine, which may provide future guidance and facilitate further research in this area. The successful journey of these analogues to clinical utility is a healthy and happy sign and an index of bright future, and we hope that this review will provide enough input to ignite the minds.

**ACKNOWLEDGEMENTS:**

Authors wish to thank Jeypore College of Pharmacy for providing library facility to carry out this review study.

**REFERENCES:**

1. Worku A, Mahmood SM. Role of taurine in the vasculature: an overview of experimental and human studies. *Am J Cardiovasc Dis*, 2011; 1(3): 293-311.
2. Schuller-Levis GB, Park E. Taurine: new implications for an old amino acid. *FEMS Microbiol Letters*, 2003; 226: 195-202.
3. Tiedemann F, Gmelin LG. Einige neue Bestandtheile der Galle des Ochsen. *Annalen der Physik*, 1827; 85 (2): 326-337.
4. Jong CJ1, Azuma J, Schaffer S. Mechanism underlying the antioxidant activity of taurine: prevention of mitochondrial oxidant production. *Amino Acids*, 2012; 42(6): 2223-2232.
5. Gupta RC, Win T, Bittner S. Taurine Analogues; A New Class of Therapeutics: Retrospect and Prospects. *Curr Med Chem*, 2005; 12(17): 75-82.
6. Huxtable RJ. Physiological actions of taurine. *Physiol Rev*, 1992; 72: 101-163.
7. Kim SJ, Ramesh C, Gupta H, Lee W. Taurine diabetes interaction: from involvement to protection. *J Biol Regul Homeost Agents*, 2007; 21: 63-77.
8. Yamori Y, Taguchi T, Hamada A, Kunimasa K, Mori H, Mori M. Taurine in health and diseases: consistent evidence from experimental and epidemiological studies. *J Biomed Sci*, 2010; 17(1): S6.

9. Gupta RC, Seki Y, Yosida J. Role of taurine in spinal cord injury. *Curr Neurovasc Res*, 2006; 3: 225-235.
10. Mozaffari MS, Borke JL. Taurine in submandibular gland of the rat: effect of muscarinic drugs. *J Histochem Cytochem*, 2002; 50: 527-532.
11. Rahman MM, Park HM, Kim SJ, Go HK, Kim GB, Hong CU, *et al.* Taurin prevents hypertension and increases exercise capacity in rats with fructose-induced hypertension. *Am J Hypertens*, 2011; 24: 574-581.
12. Nandhini AT, Thirunavukkarasu V, Anuradha CV. Taurine modifies insulin signaling enzymes in the fructose fed insulin resistant rats. *Diabetes Metab*, 2005; 31: 337-344.
13. Kuwahara M, Kawaguchi T, Ito K, Tsubone H. Effects of taurine on cardiovascular and autonomic nervous functions in cold exposed rats. *Adv Exp Med Biol*, 2009; 643: 533-540.
14. Meldrum MJ, Tu R, Patterson T, Dawson RJ, Petty T. The effect of taurine on blood pressure, and urinary sodium, potassium and calcium excretion. *Adv Exp Med Biol*, 1994; 359:207-15.
15. Abebe W, Mozaffari MS. Taurine depletion alters vascular reactivity in rats. *Can J Physiol Pharmacol*, 2003; 81: 903-909.

**Conflict of Interest:** None

**Source of Funding:** Nil

**Paper Citation:** Panda S, *et al.* A Review on Taurine - A Magic Molecule. *J Pharm Adv Res*, 2018; 1(2): 84-87.